



HERNANDO COUNTY
BOARD OF COUNTY COMMISSIONERS
DEPARTMENT OF DEVELOPMENT SERVICES
1653 BLAISE DRIVE
BROOKSVILLE, FLORIDA 34601

CU2605
WYATT HOLLY L
18185 FERRY AVE
BROOKSVILLE FL 34604-7555,

34604-7555



TAMPA FL 335
2 APR 2026 PM 6 L



quadtent
FIRST-CLASS MAIL
PRSRT - IMI
\$000.68
04/03/2026 ZIP 34601
043M32206716

US POSTAGE

Mukesh H. Mehta, M.D.
Pulmonary/Critical Care/Internal Medicine

NPI# 1689644304

DEA # BM1563382

Phone: 352-754-7222

Phone: 352-597-0690

Name Holly wyatt Age 5/3/72

Address _____ Date _____

R

7/30/26

Pt Evaluated in
office for Dyspnea.
was recently diagnosed &
confirmed Asthma
likely due to environmental
factors

Refill _____ Times _____ om

7141 Mariner Blvd. ♦ Spring Hill, FL 34609



DEPARTMENT OF PLANNING AND ZONING

PLANNING DIVISION

1653 BLAISE DRIVE ♦ BROOKSVILLE, FLORIDA 34601

P 352.754.4057 ♦ F 352.754.4420 ♦ W www.HernandoCounty.us

NOTICE OF PUBLIC HEARING HERNANDO COUNTY, FLORIDA

PLANNING AND ZONING COMMISSION

County records show you to be an owner of property within 500 feet of property that is being considered for a Conditional Use Permit, and you are hereby notified that the Planning and Zoning Commission will hold a public hearing on **April 13, 2026**, to consider the following petition. **The meeting starts at 9:00 AM and the case will be heard thereafter in the order established when the agenda is published.** Any person wishing to be heard on this matter may either be present to speak on the issue at the hearings or may provide written comments to this office. Written comments will become part of the public record.

APPLICANT: Sanchez Miledy M, Regus Rafael A, Brown Johanna

FILE NUMBER: CU-26-05

PURPOSE: Conditional Use Permit for a Second Residence Due to Medical Hardship

GENERAL

LOCATION: Southwest corner of Evening Star Avenue and Station Boulevard

PARCEL KEY

NUMBER: 203229

All hearings are held in the John Law Ayers County Commission Chambers, which is located in the Hernando County Government Center, 20 North Main Street, Room 160, Brooksville, Florida. You are further advised that if a person decides to seek review of any decision made by the board with respect to any matter considered at such meeting or hearing, he will need a record of the proceedings, and that, for such purpose, he may need to ensure that a verbatim record of the proceedings is made which record includes the testimony and evidence upon which the review is to be based.

If you should have any additional questions regarding this issue, please contact the Planning Department at planning@hernandocounty.us, 352-754-4057, or in writing at: Hernando County Planning Department, 1653 Blaise Drive, Brooksville, Florida 34601. Furthermore, information regarding this application may be obtained from the County's website at www.hernandocounty.us - follow the Board Agendas and Minutes link to the specified public hearing.

In accordance with the Americans with Disabilities Act, persons with disabilities needing a special accommodation to participate in this proceeding should contact Maryellen Waite, Hernando County Risk Management, 15470 Flight Path Drive Brooksville, FL 34604, telephone (352) 442-7787, no later than three (3) days prior to the proceedings. If hearing impaired, please call 1-800-676-3777 for assistance.

Sincerely,

A handwritten signature in purple ink, reading "Michelle Miller", is written over a faint, larger purple signature.

Michelle Miller, Planning Manager
Hernando County Planning Department

Referral Requisition

AdventHealth Medical Group Family and Internal Medicine at Parsons Park 116 PARSONS PARK DRIVE BRANDON FL 33511-6066 Phone: 813-684-5255 Fax: 813-654-7457	Date: Jan 14, 2026
---	---------------------------

Ambulatory referral to Pulmonology
Diagnosis: Recurrent respiratory infection (J98.8)

Patient Information

Legal Name: Holly Wyatt	MRN: 28296594
18185 FERRY AVE BROOKSVILLE FL 34604-7555	Phone: 352-397-8371 [REDACTED] Sex: F
Payor: MEDICARE Plan: MEDICARE A/B	Group #: Subscriber ID: 9XX0Q52DG42

Referral Information

Referred By	Referred To
Roxann Casero Rodriguez, APRN 116 Parsons Park Drive Brandon FL 33511 Phone: 813-684-5255 Fax: 813-654-7457	Diagnoses: Recurrent respiratory infection Order: Ambulatory Referral To Pulmonology Mukesh Mehta 7141 mariner blvd SPRING HILL FL 34609 Phone: 352-754-7222 Fax: 352-754-7224

Additional Referral Information:

# Visits: 1	Urgency: Routine
Start Date: Jan 14, 2026	End Date: To be determined by Insurer
Referral Type: Consultation [3]	

Electronically signed by: RODRIGUEZ, ROXANN CASERO at 11:20 am on 1/14/2026

Ordering Provider: RODRIGUEZ, ROXANN CASERO NPI: 1710327622

Authorizing Provider: RODRIGUEZ, ROXANN CASERO NPI: 1710327622

This document serves as a request of services and does not constitute Insurance authorization or approval of services. To determine eligibility, please contact the members Insurance carrier to verify and review coverage.

If you have medical questions regarding this request for services. Please contact AdventHealth Medical Group Family and Internal Medicine at Parsons Park at 813-684-5255 between the hours of 8:00am - 5:00pm (Mon-Fri).



Jan. 18th, 10:12 Am



Sunday Jan. 18th 2026 10:14 AM
Day I called the Sheriff out
But the Brown's wouldn't open
the gate for the Sheriff, made a
report.



Sunday Jan. 18th 26 10:19 Am



Sunday Jan. 18th 26
10:15am The dogs pin for restroom on fenceline
they dig and eat whatever is down there they
get very sick! I Lost one of my yorkies
she got so sick she passed away right before
court last time! 🐾



Jan. 18th, 2026
10:03 AM




Feb. 2nd / 24 8:29 AM



Look How
close they
Are to my
fence this is
at my living
Room
Area

Jan. 24th 26 9:02 AM


This is waiting like they used when I called Health
dept. back in July when the raw sewage was all
over there ground and mine! I had toilet paper
and turds in my yard was so wet it was puddling



Jan. 24, 26
3:00 pm (NO Rain this day)
OR Prior



mon. Jan. 19th 5:43pm



Feb. 1 st/₂₆ 6:29 pm



Feb. 1st/₂₆ 6:29 pm



Feb 2nd 8:20 AM
24



Feb 1st 26 8:35 AM



Feb. 1st
26 9:48 Am



Sunday Feb. 1st 26 11:38 AM



Feb. 1st 11:39 AM



Feb. 1st 26 11:39 AM



Feb. 1st 26 6:27pm

Presented During Public Comment CU-26-05

AFTER VISIT SUMMARY

Holly Wyatt MRN: 28296594
CEID: ADV-DXMV-8DMW-NJ2L

1/14/2026 9:15 AM AdventHealth Medical Group
Family and Internal Medicine at Parsons Park 813-684-5255

Instructions from Roxann Rodriguez, APRN



Today's medication changes

- ➔ **START** taking:
 - amoxicillin-clavulanate (Augmentin)
 - desvenlafaxine succinate ER (Pristiq)
 - methyLPREDNISolone (Medrol Dosepak)
- ✕ **STOP** taking:
 - cephalexin 500 MG capsule (Keflex)
 - PARoxetine 20 MG tablet (Paxil)
 - tiZANidine 4 MG capsule (Zanaflex)

Accurate as of January 14, 2026 11:26 AM.
Review your updated medication list below.



Pick up these medications at CVS 17535 IN TARGET - SPRING HILL, FL - 3101 AERIAL WAY

- amoxicillin-clavulanate 875-125 MG tablet
Your estimated payment per fill: \$6
- desvenlafaxine succinate ER 50 MG 24 hr tablet
Your estimated payment per fill: \$10
- methyLPREDNISolone 4 MG tablets
Your estimated payment per fill: \$10

Address: 3101 AERIAL WAY, SPRING HILL FL 34604
Phone: 352-796-0286



Referrals made today

Ambulatory referral to Pulmonology (Mukesh Mehta)
Expires: 1/14/2027 (requested)

X Ray Chest 2 Views
Expires: 1/14/2027 (requested)

X Ray Cervical Spine 2 - 3 Views
Expires: 1/14/2027 (requested)

X Ray Spine Lumbar 2 - 3 Views
Expires: 1/14/2027 (requested)

Today's Visit



You saw Roxann Rodriguez, APRN on Wednesday January 14, 2026. The following issues were addressed:

- Dysuria
- Polyarthralgia
- Bilateral carpal tunnel syndrome
- Chronic neck pain
- Chronic midline low back pain with bilateral sciatica
- Post menopausal syndrome
- Recurrent respiratory infection
- Mixed anxiety and depressive disorder
- Stress and adjustment reaction
- Mixed hyperlipidemia
- Environmental and seasonal allergies
- Prediabetes
- Overweight (BMI 25.0-29.9)
- Dietary counseling



Blood Pressure
120/82



BMI
26.29



Weight
158 lb



Height
5' 5"



Temperature (Touch Free)
97.9 °F



Pulse
78



Respiration
15



Oxygen Saturation
97%

Done Today

Urine culture (clean catch) for Dysuria
POC Urinalysis dipstick auto, manually resulted for Dysuria

Michelle Zering
352-754-4057
conditional use rent
20

FL. State
Before 1-800-320-0519
Call 813-472-5700
EPA 1-800-241-1754



Labs ordered today

CBC Auto Diff, Reflex Manual Diff if Indicated
Complete as directed by your provider.

Comprehensive Metabolic Panel (CMP)
Complete as directed by your provider.

Hemoglobin A1C
Complete as directed by your provider.

Lipid Panel
Complete as directed by your provider.



Follow up in about 6 weeks

(around 2/25/2026).

What's Next

FEB
25
2026

Office Visit with Roxann Rodriguez, APRN

Wednesday February 25 9:30 AM (Arrive by 9:15 AM)

Please bring any insurance information and a copayment if required
by your insurance company.

AdventHealth Medical Group Family and
Internal Medicine at Parsons Park
116 PARSONS PARK DRIVE
BRANDON FL 33511-6066
813-684-5255

You are allergic to the following

Allergen

Reactions

Duloxetine

Vomiting

Sulfa Antibiotics

Hives

Any sulfa drugs

Thank you for choosing AdventHealth for your healthcare needs. We encourage you to prioritize your annual screenings and regular check-ups. These routine visits are vital for early detection and prevention of potential health issues, ensuring you stay on track for a healthier future. Remember, proactive care today leads to a better quality of life tomorrow. Please reach out to your Primary Care team to discuss being scheduled for any outstanding items.

AdventHealth

To connect and view your health records, send messages to your doctor, renew your prescriptions, schedule appointments, and more, create an account by downloading our app or going to <https://account.adventhealth.com/register/>

Your Medication List as of January 14, 2026 11:26 AM

 Always use your most recent med list.

	acyclovir 5 % cream Commonly known as: Zovirax	Apply topically 5 (five) times a day.
 START	amoxicillin-clavulanate 875-125 MG tablet Commonly known as: Augmentin Started by: Roxann Rodriguez, APRN	Take 1 tablet by mouth in the morning and 1 tablet before bedtime. Do all this for 7 days.
	azelastine 0.1 % nasal spray Commonly known as: Astelin	SPRAY 1 SPRAY INTO EACH NOSTRIL IN THE MORNING AND BEFORE BEDTIME AS DIRECTED
 START	desvenlafaxine succinate ER 50 MG 24 hr tablet Commonly known as: Pristiq Started by: Roxann Rodriguez, APRN	Take 1 tablet (50 mg total) by mouth 1 (one) time each day. Do not crush, chew, or split.
	estradiol 2 MG tablet Commonly known as: Estrace	TAKE 1 TABLET BY MOUTH EVERY DAY
	fluticasone 50 MCG/ACT nasal spray Commonly known as: Flonase	INSTILL 2 SPRAYS PER NOSTRIL ONCE DAILY
	HYDROcodone-acetaminophen 10-325 MG tablet Commonly known as: Norco	Take 1 tablet by mouth in the morning and 1 tablet at noon and 1 tablet before bedtime.
	meloxicam 15 MG tablet Commonly known as: Mobic	Take 1 tablet (15 mg total) by mouth 1 (one) time each day.
 START	methyLPREDNISolone 4 MG tablets Commonly known as: Medrol Dosepak Started by: Roxann Rodriguez, APRN	Take as directed on package.
	nalOXone 4 mg/0.1 mL nasal spray Commonly known as: Narcan	USE SPRAY AS DIRECTED
	simvastatin 40 MG tablet Commonly known as: Zocor	TAKE 1 TABLET BY MOUTH EVERY DAY IN THE EVENING

Your Results

POC URINALYSIS DIPSTICK AUTO - PENDING

Collected on 1/14/2026

Authorized by Roxann Rodriguez, APRN

Spiritual Care Hotline

Phone: 833-258-2414

Our Spiritual Care Hotline is a confidential, anonymous, and free resource for AdventHealth patients, team members and community members. Our team is available:

- Monday - Thursday: 9:30am - 5:30pm EST
- Friday 9:30am to 3pm EST

My Chart Announcements

We now release results and clinical notes to you as soon as they are available via MyChart so you may see results before your healthcare provider has seen them. Please allow your provider time to review the results and follow-up with you. For outpatient results, if you don't hear from your provider in 3 business days, or if you have an immediate concern, please send your provider a message via MyChart.

Medication Guide
Montelukast Sodium (mon te loo' kast soe' dee um)
Tablets USP, and
Chewable Tablets

What is the most important information I should know about montelukast sodium?

Serious mental health problems have happened in people taking montelukast sodium or even after treatment has stopped. This can happen in people with or without a history of mental health problems. Stop taking montelukast sodium and tell your healthcare provider right away if you or your child have any unusual changes in behavior or thinking, including any of these symptoms:

- agitation, including aggressive behavior or hostility
- attention problems
- bad or vivid dreams
- depression
- disorientation (confusion)
- feeling anxious
- irritability
- hallucinations (seeing or hearing things that are not really there)
- memory problems
- obsessive-compulsive symptoms
- restlessness
- sleep walking
- stuttering
- suicidal thoughts and actions (including suicide)
- tremor
- trouble sleeping
- uncontrolled muscle movements

What is Montelukast sodium?

Montelukast sodium is a prescription medicine that blocks substances in the body called leukotrienes. This may help to improve symptoms of asthma and inflammation of the lining of the nose (allergic rhinitis). Montelukast sodium does not contain a steroid.

Montelukast sodium is used to:

1. Prevent asthma attacks and for the long-term treatment of asthma in adults and children ages 2 years and older. **Do not take montelukast sodium if you need relief right away for a sudden asthma attack.** If you have an asthma attack, you should follow the instructions your healthcare provider gave you for treating asthma attacks.
2. Prevent exercise-induced asthma in people 6 years of age and older.
3. Help control the symptoms of allergic rhinitis such as sneezing, stuffy nose, runny nose, and itching of the nose. Montelukast sodium is used to treat the following in people who have already taken other medicines that did not work well enough or in people who could not tolerate other medicines:

- outdoor allergies that happen part of the year (seasonal allergic rhinitis) in adults and children ages 2 years and older, **and**
- indoor allergies that happen all year (perennial allergic rhinitis) in adults and children ages 2 years and older.

Do not take Montelukast sodium if you are allergic to any of its ingredients. See the end of this Medication Guide for a complete list of the ingredients in montelukast sodium.

Before taking montelukast sodium, tell your healthcare provider about all your medical conditions, including if you:

- are allergic to aspirin.
- have phenylketonuria. Montelukast sodium chewable tablets contain aspartame, a source of phenylalanine.
- have or have had mental health problems.
- are pregnant or plan to become pregnant. Talk to your healthcare provider if you are pregnant or plan to become pregnant, montelukast sodium may not be right for you.
- are breastfeeding or plan to breastfeed. It is not known if montelukast sodium passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking montelukast sodium.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how montelukast sodium works, or montelukast sodium may affect how your other medicines work.

How should I take montelukast sodium?

For anyone who takes montelukast sodium:

- Take montelukast sodium exactly as prescribed by your healthcare provider. Your healthcare provider will tell you how much montelukast sodium to take, and **when to take it**.
- Stop taking montelukast sodium and tell your healthcare provider right away if you or your child have any unusual changes in behavior or thinking.
- You can take montelukast sodium with food or without food.
- **If you or your child misses a dose of montelukast sodium, just take the next dose at your regular time.** Do not take 2 doses at the same time.
- If you take too much montelukast sodium, call your healthcare provider right away.

For adults and children 2 years of age and older with asthma:

- Take montelukast sodium 1 time each day, in the evening. Continue to take montelukast sodium every day for as long as your healthcare provider prescribes it, even if you have no asthma symptoms.
- Tell your healthcare provider right away if your asthma symptoms get worse, or if you need to use your rescue inhaler medicine more often for asthma attacks.
- Always have your rescue inhaler medicine with you for asthma attacks.
- Continue to take your other asthma medicines as prescribed unless your healthcare provider tells you to change how you take these medicines.

For people 6 years of age and older for the prevention of exercise-induced asthma:

- Take montelukast sodium at least 2 hours before exercise.
- Always have your rescue inhaler medicine with you for asthma attacks.
- If you take montelukast sodium every day for chronic asthma or allergic rhinitis, **do not** take another dose to prevent exercise-induced asthma. Talk to your healthcare provider about your treatment for exercise-induced asthma.
- **Do not take 2 doses of montelukast sodium within 24 hours (1 day).**

For anyone 2 years of age and older with seasonal allergic rhinitis, or for anyone 2 years of age and older with perennial allergic rhinitis:

- Take montelukast sodium 1 time each day, at about the same time each day.

What should I avoid while taking montelukast sodium?

If you have asthma and aspirin makes your asthma symptoms worse, continue to avoid taking aspirin or other medicines called non-steroidal anti-inflammatory drugs (NSAIDs) while taking montelukast sodium.

What are the possible side effects of montelukast sodium?

Montelukast sodium may cause serious side effects, including:

- See “**What is the most important information I should know about montelukast sodium?**”
- **Increase in certain white blood cells (eosinophils) and possible inflamed blood vessels throughout the body (systemic vasculitis).** Rarely, this can happen in people with asthma who take montelukast sodium. This sometimes happens in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered.

Tell your healthcare provider right away if you get one or more of these symptoms:

- a feeling of pins and needles or numbness of arms or legs
- a flu-like illness
- rash
- severe inflammation (pain and swelling) of the sinuses (sinusitis)

The most common side effects of montelukast sodium include:

- upper respiratory infection
- fever
- headache
- sore throat
- cough
- stomach pain
- diarrhea
- earache or ear infection
- flu

- runny nose
- sinus infection

These are not all the possible side effects of montelukast sodium. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store montelukast sodium?

- Store montelukast sodium at room temperature between 20°C to 25° C (68°F to 77° F).
- Keep montelukast sodium in the container it comes in.
- Keep montelukast sodium in a dry place and away from light.
- **Keep montelukast sodium and all medicines out of reach of children.**

General Information about the safe and effective use of montelukast sodium

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use montelukast sodium for a condition for which it was not prescribed. Do not give montelukast sodium to other people even if they have the same symptoms you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about montelukast sodium that is written for health professionals.

What are the ingredients in montelukast sodium?

Active ingredient: montelukast sodium

Inactive ingredients:

- 10 mg tablets: croscarmellose sodium, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate and microcrystalline cellulose. The film coating consists of: hypromellose, iron oxide yellow, iron oxide red, iron oxide black, polyethylene glycol 400 and titanium dioxide.
- 4 mg and 5 mg chewable tablets: aspartame, cherry flavor, croscarmellose sodium, hydroxypropyl cellulose, mannitol, magnesium stearate, microcrystalline cellulose and red ferric oxide.

People with Phenylketonuria: Montelukast sodium 4 mg (montelukast) chewable tablets contain 0.6 mg of phenylalanine and montelukast sodium 5 mg (montelukast) chewable tablets contain 0.75 mg of phenylalanine.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Rx only

Distributor:

Dr. Reddy's Laboratories Inc.,

Princeton, NJ 08540

Made in India

Revised: 10/2022

Dispense with Medication Guide available at: www.drreddys.com/pi/montelukasttabsandchewtabs.pdf

WAITING
WY 31

Promised: 1/27/26, 4:01 PM

Scripts: 01



Wyatt, Holly L

18185 Ferry Ave, Brooksville, FL 34604

TEL: (352) 397-8371

Prescription Information

 MORNING	SEE DIRECTIONS	HYDROCORTISONE 10 MG TABLET Common brand(s): Cortef Take 1 tablet by mouth daily
 MIDDAY		
 EVENING		
 BEDTIME		
▲ PHARMACY ADVICE See back for more information		Important Information - Take this medication with food. - Take or use this exactly as directed. Do not skip doses or discontinue.

Receipt & Refill Information

CVS Pharmacy 3101 Aerial Way Brooksville, FL	STORE#: 17535	HYDROCORTISONE 10 MG TABLET
STORE TEL: (352) 796-0286 RX: 190174 00		NDC: 64380-0971-06 DAW: 0 QTY: 7 EA
		CAP: Safety MFR PKG: Yes
		REFILL: 0 Refills MFR: STRIDES PHARMA PRSCBR: Mukesh Mehta DAYS SUPPLY: 7 DATE FILLED: 1/27/26
RETAIL PRICE: \$14.99		AMOUNT DUE: \$1.90

Notes from the Pharmacy

	Vaccines you may be due for: -Flu -Covid -RSV View back of the label for more! Scan the QR to schedule now!
--	---



DRUG STORAGE TYPE: 13

RX 190174 00
DATE: 1/27/26 TP: 22845

HYDROCORTISONE 10 MG TABLET

WYATT, HOLLY L
RX 190174 00

WYATT, HOLLY L
18185 FERRY AVE
BROOKSVILLE, FL 34604

MEHTA, M
7141 MAINER BLVD
SPRING HILL, FL
TEL: (352) 597-0690

HYDROCORTISONE 10 MG TABLET

MANUFACTURER: STRIDES PHARMA

This is a WHITE, OVAL-shaped TABLET imprinted with H10 on the front.

ACTIVE INGREDIENT:

HYDROCORTISONE - ORAL - (HYE-droe-KOR-ti-sone)

COMMON BRAND NAME(S):

Cortef

USES:

Hydrocortisone is a man-made version of a natural substance (cortisol) made by the adrenal gland. It is used to treat conditions such as arthritis, blood/hormone/immune system disorders, skin and eye conditions, breathing problems, cancer, and severe allergies. It decreases your immune system's response to various diseases to reduce symptoms such as pain, swelling and allergic-type reactions. Hydrocortisone is also used to treat low cortisol levels caused by diseases of the adrenal gland (such as Addison's disease, adrenocortical insufficiency). Corticosteroids are needed in many ways for the body to function well. They are important for salt and water balance and keeping blood pressure normal.

HOW TO USE:

Take this medication by mouth, with food or milk to prevent stomach upset, exactly as directed by your doctor. Take this medication with a full glass of water (8 ounces/240 milliliters) unless your doctor directs you otherwise. Follow the dosing schedule carefully. The dosage and length of treatment are based on your medical condition and response to treatment. Your doctor may direct you to take hydrocortisone 1 to 4 times a day or take a single dose every other day. It may help to mark your calendar with a reminder. Do not stop taking this medication without consulting your doctor. Some conditions may become worse when this drug is suddenly stopped. Your dose may need to be gradually decreased. If you suddenly stop using this medication, you may have withdrawal symptoms (such as weakness, weight loss, nausea, muscle pain, headache, tiredness, dizziness). To help prevent withdrawal, your doctor may lower your dose slowly. Withdrawal is more likely if you have used hydrocortisone for a long time or in high doses. Tell your doctor or pharmacist right away if you have withdrawal. See also Precautions section. Tell your doctor if you have any other health conditions or are taking other medications. Tell your doctor if you are pregnant or planning to get pregnant, as this medication may cause complications. Tell your doctor if you are breastfeeding, as this medication may pass into your breast milk and affect your baby. Tell your doctor if you have any other health conditions or are taking other medications. Tell your doctor if you are pregnant or planning to get pregnant, as this medication may cause complications. Tell your doctor if you are breastfeeding, as this medication may pass into your breast milk and affect your baby.

SIDE EFFECTS:

Nausea, heartburn, headache, trouble sleeping, increased sweating, or acne may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may rarely make your blood sugar rise, which can cause or worsen diabetes. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. If you already have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet. Tell your doctor right away if you have any serious side effects, including: signs of infection (such as sore throat that doesn't go away, fever, chills), weakness, swelling ankles/feet, unusual weight gain, vision problems, easy bruising/bleeding, puffy face, mental/mood changes (such as depression, mood swings, agitation), muscle weakness/pain, thinning skin, slow wound healing, menstrual period changes, irregular heartbeat, bone pain, seizures, symptoms of stomach/intestinal bleeding (such as stomach/abdominal pain, black/tarry stools, vomit that looks like coffee grounds), red/purple

breast milk. However, this drug is unlikely to harm a nursing infant. Consult your doctor before breastfeeding.

DRUG INTERACTIONS:

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: aldesleukin, desmopressin, mifepristone, other drugs that can cause bleeding/bruising (such as antiplatelet drugs like clopidogrel, NSAIDs like ibuprofen/naproxen, "blood thinners" like warfarin/dabigatran). Aspirin can increase the risk of bleeding when used with this medication. However, if your doctor has told you to take low-dose aspirin to prevent heart attack or stroke (usually 81-162 milligrams a day), you should keep taking the aspirin unless your doctor tells you not to. Ask your doctor or pharmacist for more details. This medication may interfere with certain lab tests (such as skin tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

OVERDOSE:

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669.

NOTES:

Do not share this medication with others. If this medication is used for a long time, lab and/or medical tests (such as blood sugar/mineral levels, blood counts, blood pressure, bone density tests, eye exams, height/weight measurements, X-rays) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. This medication may cause bone loss (osteoporosis) when used for a long time. Lifestyle changes that may help reduce the risk of bone problems include doing weight-bearing exercise, stopping smoking, limiting alcohol, and getting enough calcium and vitamin D. Talk to your doctor for more details.

MISSED DOSE:

If you are taking this medication once daily and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. If you are taking this medication every other day, ask your doctor or pharmacist what you should do if you miss a dose.

STORAGE:

Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised September 2025. Copyright(c) 2026 First Databank, Inc.

It is important to dispose of unneeded or expired medications, including opioids, in a safe and secure fashion. Drug disposal units are available at select locations. Visit <https://safe.pharmacy/drug-disposal/> to find a convenient disposal location near you. You may also qualify to receive an at-home drug disposal solution when picking up certain prescriptions. See a pharmacy colleague for additional information.

A Simple Way to Manage your Prescriptions

CVS Pharmacy has made it easier for you to read and follow your prescription information. We use icons to show meaningful pharmacy advice on when to take each medication dose, so your prescriptions can work harder for you.

As always, your CVS Pharmacy team welcomes your questions.

Any questions about your medication? Contact your pharmacist:
Marthe Deshauteurs, RPh
TEL: (352) 796-0286

Keep Out of Reach of Children. Important Disclaimer: The side effects described in this document do not reflect all the possible risks that could be caused by this medication. For further information, please consult with your physician about the uses, precautions and risks of the medication specific to your health. This information is obtained from First DataBank for use as an educational aid.

Item#: 399469

spots on the skin. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:
Before taking hydrocortisone, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: eye disease (such as cataracts, glaucoma), heart problems (such as heart failure, recent heart attack), high blood pressure, liver disease (such as hepatitis B), kidney disease, thyroid problems, diabetes, stomach/intestinal problems (such as diverticulitis, ulcer), brittle bones (osteoporosis), current/past infections (such as tuberculosis, positive tuberculosis test, herpes, fungal), bleeding problems, blood clots, mental/mood conditions (such as psychosis, anxiety, depression), low salts in the blood (such as low potassium or calcium), seizures, a certain adrenal gland tumor (pheochromocytoma). This medication may cause stomach bleeding. Daily use of alcohol while using this medication may increase your risk for stomach bleeding. Limit alcoholic beverages. Consult your doctor or pharmacist for more information. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Using corticosteroid medications for a long time can make it more difficult for your body to respond to physical stress. Before having surgery or emergency treatment, or if you get a serious illness/injury, tell your doctor or dentist that you are using this medication or have used this medication within the past 12 months. Tell your doctor right away if you develop unusual/extreme tiredness or weight loss. If you will be using this medication for a long time, carry a warning card or medical ID bracelet that identifies your use of this medication. This medication may mask signs of infection. It can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using hydrocortisone before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). This medication may slow down a child's growth if used for a long time. Consult the doctor or pharmacist for more details. See the doctor regularly so your child's height and growth can be checked. Older adults may be more sensitive to the effects of this drug, especially bone loss/pain, stomach/intestinal bleeding, and mental/mood changes (such as confusion). During pregnancy, hydrocortisone should be used only when clearly needed. Discuss the risks and benefits with your doctor. Infants born to mothers who have been using this medication for an extended period of time may have hormone problems. Tell your doctor right away if you notice symptoms such as nausea/vomiting that doesn't stop, severe diarrhea, or weakness in your newborn. It is unknown if this drug passes into

Our records indicate the following vaccines are recommended for you

Flu

Everyone over 6 months should get a seasonal flu vaccine to help protect themselves

Covid

Recommended based on individual decision making, with most benefit for high risk individuals

RSV

Everyone 75 and older, and 50 to 74 at increased risk should receive to help prevent severe illness

Hepatitis B

Getting the Hepatitis B vaccine series is the best way to help prevent Hepatitis B infection

Scan the QR code below to schedule your vaccine!



Pharmacy Advice*

The dose schedule contains your pharmacy's recommendation on the best time to take this medication, taking into account other medications you have received from CVS Pharmacy. It does not take into account medications you are receiving from other pharmacies. This recommendation may change if you add or stop taking other medications after 1/27/26 3:49 PM. If you have questions about your medication or whether they can be taken at another time, please consult your pharmacist.

WY Rx: XXX0174 00



4301901740006438009710617535000000

Personalized Education Printout

Your pharmacy has relevant information to share with you about one or more of your medications.

Prescription being filled: HYDROCORTISONE 10 MG TABLET

In combination with: CIPROFLOXACIN HCL 500 MG TAB

- Tell your medical provider if you hear a snap or popping sound after an injury, experience rapid or immediate bruising, are unable to utilize the affected area or bear weight, or the injured area becomes deformed. These may be signs of a ruptured tendon.
- Your medicines may increase your risk of a tendon rupture. If you experience pain or inflammation, rest and refrain from exercise until your medical provider tells you that you may exercise.
- If you have questions or concerns or stop taking your medicines for any reason, contact your medical provider.

If you have any questions, please call or visit your local CVS Pharmacy:

3101 Aerial Way
Brooksville, FL
(352) 796-0286

Copyright (c) 2026 First Databank, Inc.

Private and Confidential. Intended for patient or caregiver only. If you have received this document in error, please notify CVS Pharmacy immediately.

Before having surgery or procedures such as X-rays with dye, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Children may be more sensitive to the side effects of this drug, especially slowed breathing and uncontrolled movements (see also Warning section). Children may also experience excitability rather than drowsiness. Special caution should be taken in children with a family history of sudden infant death syndrome (SIDS), those who have a severe loss of fluid (dehydration), and those who are hard to wake up from sleep. Older adults may be more sensitive to the side effects of this drug, especially drowsiness, confusion, constipation, or trouble urinating. Drowsiness and confusion can increase the risk of falling. During pregnancy, this medication should be used only if clearly needed. Discuss the risks and benefits with your doctor. Dextromethorphan passes into breast milk. It is unknown if promethazine passes into breast milk. Consult your doctor before breastfeeding.

DRUG INTERACTIONS:

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: diphenhydramine or other antihistamines applied to the skin (such as anti-itch cream, ointment, spray), metoclopramide, rolapitant. Taking certain MAO inhibitors with this medication may cause a serious (possibly fatal) drug interaction. Avoid taking isocarboxazid, metaxalone, methylene blue, moclobemide, phenelzine, procarbazine, rasagiline, safinamide, selegiline, or tranylcypromine during treatment with this medication. Most MAO inhibitors should also not be taken for two weeks before treatment with this medication. Ask your doctor when to start or stop taking this medication. The risk of serious side effects (such as slow/shallow breathing, severe drowsiness/dizziness) may be increased if this medication is taken with other products that may also cause drowsiness or breathing problems. Tell your doctor or pharmacist if you are taking other products such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or other antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely. Dextromethorphan is available in both prescription and nonprescription products. Check the labels of all your medications to make sure you are not taking more than one product containing dextromethorphan. This medication may interfere with certain lab tests (such as urine pregnancy tests, skin allergy tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

OVERDOSE:

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669. Symptoms of overdose may include: severe drowsiness, inability to wake up, slow/shallow breathing, slow heartbeat, hallucinations, seizures.

NOTES:

Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another condition unless told to do so by your doctor. A different medication may be necessary in those cases.

MISSED DOSE:

If you are prescribed this medication on a regular schedule and miss a

dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised July 2025. Copyright(c) 2026 First Databank, Inc.

It is important to dispose of unneeded or expired medications, including opioids, in a safe and secure fashion. Drug disposal units are available at select locations. Visit <https://safe.pharmacy/drug-disposal/> to find a convenient disposal location near you. You may also qualify to receive an at-home drug disposal solution when picking up certain prescriptions. See a pharmacy colleague for additional information.

A Simple Way to Manage your Prescriptions

CVS Pharmacy has made it easier for you to read and follow your prescription information. We use icons to show meaningful pharmacy advice on when to take each medication dose, so your prescriptions can work harder for you.

As always, your CVS Pharmacy team welcomes your questions.

Any questions about your medication? Contact your pharmacist:

Marthe Deshauteurs, RPh

TEL: (352) 796-0286

Keep Out of Reach of Children. Important Disclaimer: The side effects described in this document do not reflect all the possible risks that could be caused by this medication. For further information, please consult with your physician about the uses, precautions and risks of the medication specific to your health. This information is obtained from First DataBank for use as an educational aid.

Item#: 399469

pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. If you are very sleepy or have difficulty waking up after using this medication, stop taking this medication and talk to your doctor or pharmacist right away. Your dosage may need to be lower. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: decreased coordination, difficulty urinating, shaking (tremor), mental/mood changes (such as nervousness, excitement, irritability, confusion, hallucinations), abnormal/uncontrolled movements (such as fixed upward stare, neck twisting, tongue movements), slow heartbeat, fainting. Get medical help right away if you have any very serious side effects, including: slow/shallow breathing, seizures. This medication may rarely cause a very serious condition called neuroleptic malignant syndrome (NMS). Get medical help right away if you have any of the following symptoms: fever, muscle stiffness/pain/tenderness/weakness, severe tiredness, severe confusion, sweating, fast/irregular heartbeat, dark urine, signs of kidney problems (such as change in the amount of urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

See also Warning section. Before taking this medication, tell your doctor or pharmacist if you are allergic to promethazine or to dextromethorphan; or to any other phenothiazines (such as prochlorperazine); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: breathing problems (such as asthma, emphysema, sleep apnea), heart problems (such as slow/irregular heartbeat, chest pain), liver problems, difficulty urinating (such as due to an enlarged prostate), stomach/bowel problems (such as blockage, ulcer), high blood pressure, diabetes, a certain eye problem (glaucoma), seizures, blood/immune system problems (such as bone marrow depression). This drug may make you dizzy or drowsy or blur your vision. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Children should be supervised during bicycle riding and other possibly hazardous activities to avoid injury. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness. Liquid forms of this medication may contain sugar, alcohol, or aspartame. Caution is advised if you have diabetes, liver disease, phenylketonuria (PKU), or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely.

Our records indicate the following vaccines are recommended for you

Flu

Everyone over 6 months should get a seasonal flu vaccine to help protect themselves

Covid

Recommended based on individual decision making, with most benefit for high risk individuals

RSV

Everyone 75 and older, and 50 to 74 at increased risk should receive to help prevent severe illness

Hepatitis B

Getting the Hepatitis B vaccine series is the best way to help prevent Hepatitis B infection

Scan the QR code below to schedule your vaccine!



Pharmacy Advice*

The dose schedule contains your pharmacy's recommendation on the best time to take this medication, taking into account other medications you have received from CVS Pharmacy. It does not take into account medications you are receiving from other pharmacies. This recommendation may change if you add or stop taking other medications after 1/27/26 3:50 PM. If you have questions about your medication or whether they can be taken at another time, please consult your pharmacist.

ACUTE
WY

31

Promised: 1/27/26, 4:10 PM

Scripts: 01

27 0190173 000 000 00 0001000

Wyatt, Holly L

18185 Ferry Ave, Brooksville, FL 34604

TEL: (352) 397-8371

Prescription Information

MORNING

MIDDAY

EVENING

BEDTIME

SEE DIRECTIONS

PROMETHAZINE-DM

6.25-15 MG/5ML

Take 5 mL by mouth 3 times a day

Important Information

- May cause drowsiness. Use care when operating a vehicle, vessel or machine.

- This medication may cause dizziness.

- Avoid prolonged exposure to direct and/or artificial sunlight.

- May cause blurred vision.

- Date opened _____.

▲ PHARMACY ADVICE*

See back for more information

Receipt & Refill Information

CVS Pharmacy

3101 Aerial Way
Brooksville, FL

STORE#: 17535

STORE TEL: (352) 796-0286

RX: 190173 00

AUTH#: 260275311214375999

REFILL: 0 Refills

MFR: ANI PHARMACEUTI

PRSCBR: Mukesh Mehta

DAYS SUPPLY: 15

DATE FILLED: 1/27/26

AMOUNT DUE: \$10.00

RETAIL PRICE:\$26.09

Notes from the Pharmacy

Vaccines you may be due for:

-Flu

-Covid

-RSV

View back of the label for more!

Scan the QR to schedule now!

WYATT, HOLLY L

18185 FERRY AVE
BROOKSVILLE, FL 34604

RX 190173 00

WYATT, HOLLY L

RX 190173 00

PROMETHAZINE-DM 6.25-15 MG/5ML

RX 190173 00

DATE: 1/27/26

TP: 22845

DRUG STORAGE TYPE: 27

MEHTA, M.

7141 MARINER BLVD
SPRING HILL, FL

TEL: (352) 597-0690

PROMETHAZINE-DM 6.25-15 MG/5ML

MANUFACTURER: ANI PHARMACEUTI

This is a YELLOW SYRUP.

ACTIVE INGREDIENT:
PROMETHAZINE/DEXTROMETHORPHAN - ORAL -
(pro-METH-uh-zeen/dex-trow-meth-OR-fan)

WARNING:
Promethazine should not be used by children younger than 2 years since i might cause serious (possibly fatal) slow/shallow breathing. This problem has occurred even with normal doses in this age group. The lowest effective dosage should be used, and other drugs that affect breathing should be avoided. Get medical help right away in the unlikely event that slow/shallow breathing occurs. Promethazine should not be used by children with liver disease (including possible Reyes syndrome).

USES:
This combination medication is used to temporarily treat runny nose, coughing, and sneezing caused by the common cold, allergies, hay fever, and other breathing illnesses. Promethazine is a phenothiazine that works like an antihistamine. It blocks the effects of a certain natural substance (histamine), which causes allergy symptoms. It also has a drying effect. Dextromethorphan belongs to a class of drugs known as cough suppressants. It acts on a part of the brain (cough center) to reduce the need to cough. This product is usually not used for ongoing cough from smoking or long-term breathing problems (such as chronic bronchitis, emphysema) unless directed by your doctor. Cough-and-cold products have not been shown to be safe or effective in children younger than 6 years. Do not use this product to treat cold symptoms in children younger than 6 years unless specifically directed by the doctor. Some products (such as long-acting tablets/capsules) are not recommended for use in children younger than 12 years. Ask your doctor or pharmacist for more details about using your product safely. These products do not cure or shorten the length of the common cold and may cause serious side effects. To decrease the risk for serious side effects, carefully follow all dosage directions. Do not use this product to make a child sleepy. Do not give other cough-and-cold medication that might contain the same active ingredients (see also Drug Interactions section). Ask the doctor or pharmacist about other ways to relieve cough and cold symptoms (such as drinking enough fluids, using a humidifier or saline nose drops/spray).

HOW TO USE:
Take this medication by mouth with or without food as directed by your doctor, usually every 4 to 6 hours as needed. Since dosing recommendations may vary, carefully follow your doctor's directions for taking this medication. The dosage is based on your age, medical condition, and response to treatment. Carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose. Drink plenty of fluids while you are using this medication unless otherwise directed by your doctor. The fluid will help loosen the mucus in your lungs. Improper use of this medication (abuse) may result in serious harm (such as brain damage, seizure, death). Do not increase your dose, take it more often, or use it for a longer time than directed. Tell your doctor if your cough lasts, returns, or gets worse after 7 days or if it is accompanied by headache that doesn't go away, fever, severe sore throat, or rash.

SIDE EFFECTS:
Drowsiness, dizziness, blurred vision, nausea, constipation, or dry mouth may occur. If any of these effects last or get worse, notify your doctor or

details.

MISSED DOSE:
If you miss a dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Do not take more than 1 dose in 24 hours.

STORAGE:
Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised November 2024. Copyright(c) 2026 First Databank, Inc.

It is important to dispose of unneeded or expired medications, including opioids, in a safe and secure fashion. Drug disposal units are available at select locations. Visit <https://safe.pharmacy/drug-disposal/> to find a convenient disposal location near you. You may also qualify to receive an at-home drug disposal solution when picking up certain prescriptions. See a pharmacy colleague for additional information.

A Simple Way to Manage your Prescriptions

CVS Pharmacy has made it easier for you to read and follow your prescription information. We use icons to show meaningful pharmacy advice on when to take each medication dose, so your prescriptions can work harder for you.

As always, your CVS Pharmacy team welcomes your questions.

Any questions about your medication? Contact your pharmacist:
Neil Drouillard, RPh
TEL: (352) 796-0286

Keep Out of Reach of Children. Important Disclaimer: The side effects described in this document do not reflect all the possible risks that could be caused by this medication. For further information, please consult with your physician about the uses, precautions and risks of the medication specific to your health. This information is obtained from First DataBank for use as an educational aid.
Item#: 399469

WAITING
WY 48

Promised: 1/30/26, 3:14 PM
Scripts: 01



27 0190552 000 000 00000500

Wyatt, Holly L

18185 Ferry Ave, Brooksville, FL 34604
TEL: (352) 397-8371

Prescription Information

 MORNING	SEE DIRECTIONS	MONTELUKAST SOD 10 MG TABLET Substituted for: Singulair 10 Mg Tablet Take 1 tablet every day
 MIDDAY		
 EVENING		
 BEDTIME		
PHARMACY ADVICE: See back for more information		Important Information - Call doctor if you experience mood changes, sadness, depression or fear.

Receipt & Refill Information

CVS Pharmacy 3101 Aerial Way Brooksville, FL	STORE#: 17535	MONTELUKAST SOD 10 MG TABLET
STORE TEL: (352) 796-0286 RX: 190552 00		NDC: 55111-0725-90 DAW: 0 QTY: 90 EA
		CAP: Safety MFR PKG: Yes
		REFILL: 0 Refills MFR: DR.REDDY'S LAB PRSCBR: Mukesh Mehta DAYS SUPPLY: 90 DATE FILLED: 1/30/26
RETAIL PRICE:\$279.99		AMOUNT DUE: \$5.00

Notes from the Pharmacy



OPEN HERE

DILIG STORAGE TYPE: FAST MOVER

RX 190552 00
DATE 1/30/26 TP: 22845

MONTELUKAST SOD 10 MG TABLET

RX 190552 00

WYATT, HOLLY L

8185 FERRY AVE
BROOKSVILLE, FL 34604
7441 MARINER BLVD
SPRING HILL, FL
TEL: (352) 597-0690

WYATT, HOLLY L

MEHTA, M.

MONTELUKAST SOD 10 MG TABLET

MANUFACTURER: DR.REDDY'S LAB

This is a BROWN, SQUARE (ROUNDED CORNERS)-shaped TABLET imprinted with R on the front and 725 on the back.

ACTIVE INGREDIENT:
MONTELUKAST - ORAL - (mon-teh-LEW-kast)
COMMON BRAND NAME(S):
Singulair
WARNING:

This medication may rarely cause serious mental/mood changes. These effects may occur even after the medication has been stopped. Tell your doctor right away if you or a caregiver/family member notice symptoms such as agitation, anxiety, abnormal dreams, depression, memory/attention problems, hallucinations, trouble sleeping, or changes in behavior (such as aggression, hostility, restlessness, sleepwalking, suicidal thoughts/attempts). Talk to your doctor about the risks and benefits of this medication.

USES:
Montelukast is used to control and prevent symptoms caused by asthma (such as wheezing and shortness of breath). It is also used before exercise to prevent breathing problems during exercise (bronchospasm). This medication can help decrease the number of times you need to use your quick relief inhaler. Montelukast is also used to relieve symptoms of hay fever and allergic rhinitis (such as sneezing, stuffy/runny/itchy nose). Since there are other allergy medications that may be safer (see also Warning section), this medication should be used for this condition only when you cannot take other allergy medications or they do not work well. This medication must be used regularly to be effective. It does not work right away and should not be used to relieve sudden asthma attacks or breathing problems. If an asthma attack or sudden shortness of breath occurs, use your quick-relief inhaler as prescribed. This drug works by blocking certain natural substances (leukotrienes) that may cause or worsen asthma and allergies. It helps make breathing easier by reducing swelling (inflammation) in the airways.

HOW TO USE:
Read the Medication Guide provided by your pharmacist before you start taking montelukast and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor. The dosage is based on your age and medical condition. If you are using the chewable tablets, chew them thoroughly before swallowing. If your child cannot safely chew and swallow them, consult the doctor or pharmacist for advice. Take this medication at the same time each day. If you are taking this medication for asthma or for both asthma and allergies, take your dose in the evening. If you are taking montelukast to prevent only allergies, take your dose either in the morning or the evening. If you are taking this medication to prevent breathing problems during exercise, take your dose at least 2 hours before exercise. Do not take more than one dose in 24 hours. Do not take a dose before exercise if you are already taking this medication daily for asthma or allergies. Doing so may increase the risk of side effects. Do not increase or decrease your dose or stop using this medication without consulting your doctor. Continue to use this medication regularly to keep your asthma under control, even during sudden asthma attacks or periods when you have no asthma symptoms. Continue to also take other medications for asthma as directed by your doctor. This medication works over time and is not meant to relieve sudden attacks of asthma. If an asthma attack or other breathing problem occurs, use your quick-relief inhaler as prescribed. You

should always have a quick-relief inhaler with you. Consult your doctor or pharmacist for more details. Get medical help right away if your asthma symptoms get worse and your quick-relief inhaler is not helping. Tell your doctor promptly if asthma symptoms, breathing problems, allergy symptoms, number of times you use your rescue inhaler last or get worse.

SIDE EFFECTS:
See also Warning section. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: numbness/tingling/shooting pain in the arms or legs, sinus pain/swelling, muscle weakness, uncontrolled muscle movements, stuttering. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US - Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:
Before taking montelukast, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this drug, tell your doctor or pharmacist your medical history, especially of: liver disease, mental/mood problems (such as anxiety, depression, thoughts of suicide). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). The chewable tablets may contain aspartame. If you have phenylketonuria (PKU) or any other condition that requires you to limit/avoid aspartame (or phenylalanine) in your diet, ask your doctor or pharmacist about using this medication safely. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication passes into breast milk. Consult your doctor before breastfeeding.

DRUG INTERACTIONS:
Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

OVERDOSE:
If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669. Symptoms of overdose may include: thirst, drowsiness, inability to keep still, vomiting, or severe stomach pain.

NOTES:
Do not share this medication with others. Lab and/or medical tests (such as lung function) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more

CONTINUE READING ON THE OTHER SIDE

CVS/pharmacy

The Pharmacy Team has information to share with you, please read below.

Message Center
We have successfully processed this Rx for a 3 month supply

Employee: Neil D.



Pharmacy Advice*

The dose schedule contains your pharmacy's recommendation on the best time to take this medication, taking into account other medications you have received from CVS Pharmacy. It does not take into account medications you are receiving from other pharmacies. This recommendation may change if you add or stop taking other medications after 1/30/26 3:00 PM. If you have questions about your medication or whether they can be taken at another time, please consult your pharmacist.